

The Longitudinal and Transverse Response of the ${}^4\text{He}(e, e'p)$ Reaction in the Dip Region

A. Kozlov¹, K.A. Aniol⁵, P. Bartsch², D. Baumann², W. Bertozzi³, R. Böhm², K. Bohinc¹⁰, J.P. Chen⁴, D. Dale⁶, L. Dennis⁸, S. Derber², M. Ding², M.O. Distler², A. Dooley⁸, P. Dragovitsch⁸, M.B. Epstein⁵, I. Ewald², K.G. Fissum³, R.E.J. Florizone³, J. Friedrich², J.M. Friedrich², R. Geiges², S. Gilad³, P. Jennewein², M. Kahrau², M. Kohl⁷, K.W. Krygier², A. Liesenfeld², D.J. Margaziotis⁵, H. Merkel², P. Merle², U. Müller², R. Neuhausen², T. Pospischil², G. Riccardi⁸, R. Roche⁸, G. Rosner², D. Rowntree³, A.J. Sarty⁸, H. Schmieden², S. Širca¹⁰, J.A. Templon⁹, M. Thompson¹, A. Wagner², Th. Walcher², M. Weis², J. Zhao³, Z. Zhou³

1 – School of Physics, The University of Melbourne, Parkville 3052, VIC, Australia;

2 – Institut für Kernphysik, Universität Mainz, D-55099 Mainz, Germany;

3 – Laboratory for Nuclear Science, MIT, Cambridge, MA 02139, USA;

4 – TJNAF, Newport News, VA, USA;

5 – Department of Physics and Astr., California St. U., Los Angeles, CA 90032, USA;

6 – Department of Physics and Astr., U. of Kentucky, Lexington, KY 40506, USA;

7 – Institut für Kernphysik, Technische U. Darmstadt, D-64289 Darmstadt, Germany;

8 – Department of Physics, Florida State University, Tallahassee, FL 32306, USA;

9 – Department of Physics and Astronomy, U. of Georgia, Athens, GA 30602, USA;

10 – Institute "Jožef Stefan", University of Ljubljana, SI-1001 Ljubljana, Slovenija;

A high-resolution study of the $(e, e'p)$ reaction on ${}^4\text{He}$ was carried out at the Institut für Kernphysik in Mainz, Germany. The high quality 100 % duty factor electron beam, and the high-resolution three-spectrometer-system of the A1 collaboration were used. The measurements were done in parallel kinematics at a central momentum transfer $|\vec{q}| = 685 \text{ MeV}/c$, and at a central energy transfer $\omega = 334 \text{ MeV}$, corresponding to a value of the y -scaling variable of $+140 \text{ MeV}/c$. In order to enable the Rosenbluth separation of the longitudinal σ_L and transverse σ_T response functions (as defined in [1]), three measurements at different incident beam energies, corresponding to three values of the virtual photon polarization ϵ , were performed.

The absolute $(e, e'p)$ cross section for ${}^4\text{He}$ was obtained as a function of missing energy and missing momentum. A distorted spectral function, $S^{dist}(E_m, p_m)$ was extracted from the data using the *cc1* prescription for the elementary off-shell e - p cross section (see ref.[2]), where

$$S^{dist}(p_m, E_m) = \frac{1}{p_p^2 \sigma_{ep}} \frac{d^6 \sigma}{d\Omega_e d\Omega_p dp_e dp_p} \quad (1)$$

For the two-body breakup channel the experimental results were compared to the theoretical calculations performed by Schiavilla *et al.* [3] and Forest *et al.* [4], and to the earlier experimental momentum distributions measured at NIKHEF by van den Brand *et al.* [5] and the new results from MAMI by Florizone [6] as shown in Fig. 1. For the continuum channel, recent calculations for the ${}^4\text{He}$ spectral function by Efros *et al.* [7] are presented in Fig. 2,B to compare with the experimental results.

A Rosenbluth separation was performed for both the two-body breakup and for continuum channels, and preliminary results were obtained. The ratio σ_L/σ_T was determined and compared with predictions (see Figure 2).




Figure 1: The $p - t$ momentum distributions

The measurements show no significant strength corresponding to the $(e, e'p)$ reaction channel for missing energy values $E_m \geq 50 \text{ MeV}$ as shown in Fig. 2.

References

1. S.Boffi *et al.*, Nucl. Phys. **A435** (1985) 697
2. T. de Forest, Nucl. Phys., **A392**, 232 (1983)
3. R. Schiavilla *et al.*, Nucl. Phys. **A449** 219 (1986)
4. J.L. Forest *et al.*, Phys. Rev **C54**, 646 (1996)
5. J.F.J. van den Brand *et al.*, Nucl. Phys. **A534** (1991) 637
6. R.E.J. Florizone *et al.*, to be published in Phys. Rev. **C**
7. V. D. Efros, W. Leidemann, G. Orlandini, Phys. Rev. **C58** (1998) 582




Figure 2: Preliminary results shown as a function of missing energy. (A) Six-fold differential cross section for the three values of the virtual-photon polarization ϵ in the missing momentum range from 115 to 155 MeV/c. (B) Spectral function. (C) The ratio σ_L/σ_T (both the 2-body and continuum channels).